

QUANTIZATION OF THREE-WAVE EQUATIONS

S. SERGEEV

ABSTRACT. The subject of this paper is the consecutive procedure of discretization and quantization of two similar classical integrable systems in three-dimensional space-time: the standard three-wave equations and less known modified three-wave equations. The quantized systems in discrete space-time may be understood as the regularized integrable quantum field theories. Integrability of the theories, and in particular the quantum tetrahedron equations for vertex operators, follow from the quantum auxiliary linear problems. Principal object of the lattice field theories is the Heisenberg discrete time evolution operator constructed with the help of vertex operators.

INTRODUCTION

The three-wave equations is the example of a completely integrable classical system in $2 + 1$ dimensional space-time. These equations apply to many physical systems. The aim of this paper is the formulation of an integrable quantum field theory corresponding to the three-wave equations.

Given a classical theory with action $S[\phi]$, there is the universal prescription for the quantization. The amplitude between a state $\phi_{in}(\mathbf{r})$ at time t_1 and a state $\phi_{out}(\mathbf{r})$ at time t_2 is

$$(1) \quad \langle \phi_{out} | \phi_{in} \rangle = \int_{\substack{\phi(\mathbf{r}, t_1) = \phi_{in}(\mathbf{r}) \\ \phi(\mathbf{r}, t_2) = \phi_{out}(\mathbf{r})}} \mathcal{D}\phi \, e^{\frac{i}{\hbar} S[\phi]}$$

The perturbation-theory approach to the definition of $\mathcal{D}\phi$ we reject from every beginning. The practical way to define the measure in the Feynman integral is the discretization. It is to be understood as

$$(2) \quad \mathcal{D}\phi = \lim \prod_{i,j} d\phi(\mathbf{r}_i, t_j),$$

where the limit symbol stands for infinitely dense discretization of the space-time $(\mathbf{r}, t)$.

Thus, the first step toward the field theory is the discretization of classical system. Only then we may look for a self-consistent Heisenberg quantum mechanics on the lattice and regard it as the regularized field theory.

A schematic outlook of milestones of our method is the following. A classical theory is defined by the time dynamics for a field A , $\frac{d}{dt}A = f[A]$ (space-like degrees of freedom are omitted for brevity). The dynamics is generated by a Hamiltonian, $f[A] = \{H, A\}$, where $\{, \}$ are properly defined Poisson brackets. Corresponding discrete time dynamics is the evolution transformation $A(t + \Delta t) = F[A(t)]$. For brevity, we choose the scale $\Delta t = 1$. The key point is that making the discretization, we have

1991 *Mathematics Subject Classification.* 37K15.

Key words and phrases. Three-Wave Equations, Integrable Systems, Discrete Time Evolution Operator, Tetrahedron Equation.

to take care of the Hamiltonian structure of the dynamics. The discrete time evolution must be a canonical transformation, it should preserve properly defined Poisson algebra of observable fields on space-like lattice. The quantum algebra of observables is a result of Dirac quantization of the Poisson algebra. The quantum evolution map $A(t) \rightarrow A(t+1)$ must be the an automorphism of the algebra of observables allowing one to define the Heisenberg evolution operator,

$$(3) \quad A(t+1) = U A(t) U^{-1} .$$

Such scheme is the algebraic realization of the discrete measure (2). Operator U acts in a representation space of the algebra of observables, the amplitude (1) in the Heisenberg form is

$$(4) \quad \langle \phi_{out} | U^T | \phi_{in} \rangle = \int_{\substack{\phi_0 = \phi_{in} \\ \phi_T = \phi_{out}}} \underbrace{\prod_{t=1}^{T-1} d\phi_t}_{\mathcal{D}\phi} \underbrace{\prod_{t=1}^T \langle \phi_t | U | \phi_{t-1} \rangle}_{e^{\frac{i}{\hbar} S[\phi]}} .$$

The discretization of the space-time, construction of the quantum algebra of observables and the basis-invariant definition of Heisenberg evolution operator for the standard and modified three-wave systems is the subject of this paper.

1. THREE-WAVE EQUATIONS

We commence with a short reminding of the classical three-wave equations in three-dimensional space. In what follows, we use the short notations for the indices,

$$(5) \quad (i, j, k) = \text{any permutation of } (1, 2, 3) .$$

1.1. Standard three-wave equations. Linear problem for the standard three-wave equations is the set of six differential relations

$$(6) \quad \partial_i \psi_j = A_{ij} \psi_i , \quad i \neq j$$

for three auxiliary fields ψ_i . Consistency of (6) gives the equations for the six fields A_{ij} :

$$(7) \quad \partial_i A_{jk} = A_{ji} A_{ik} .$$

These are the equations of motion for tree-wave resonant system (see e.g. [1, 2]), or the three-wave equations for the shortness.

1.2. Modified three-wave equations. There is another type auxiliary linear problem [3], the second order differential equations for the scalar auxiliary field ϕ :

$$(8) \quad (\partial_i \partial_j - A_{ij} \partial_j - A_{ji} \partial_i + A_{ij} A_{ji}) \phi = 0 , \quad i \neq j .$$

Consistency of (8) gives similar equations for the fields A_{ij} ,

$$(9) \quad \partial_i A_{jk} = (A_{ij} - A_{ik})(A_{jk} - A_{ji}) .$$

We call them the modified three-wave equations.

1.3. Hamiltonians. It is convenient to use the single alphabetical indices instead of the numerical pairs. In this paper we will use the following convention for both systems:

$$(10) \quad \begin{aligned} A_{12} &= A_a, & A_{13} &= A_b, & A_{23} &= A_c, \\ A_{21} &= A_a^*, & A_{31} &= A_b^*, & A_{32} &= A_c^*, \end{aligned}$$

Note, our notations are not cyclic with respect to $(1, 2, 3)$.

Equations (7) and (9) are extremum conditions for the action

$$(11) \quad \mathcal{S} = \int d^3x (A_a^* \partial_3 A_a - A_b^* \partial_2 A_b + A_c^* \partial_1 A_c - V)$$

where

$$(12) \quad V = A_a^* A_b A_c^* - A_a A_b^* A_c$$

for the standard three-wave equations (7), and

$$(13) \quad V = (A_a - A_b)(A_a^* - A_c)(A_b^* - A_c^*)$$

for the modified three-wave equations (9). For a moment, we ignore the reality conditions, the star in all these notations *does not* mean the complex conjugation.

The time derivative ∂_t and space derivatives ∂_x, ∂_y for action (11) may be chosen by

$$(14) \quad \partial_1 = \partial_t - \partial_x, \quad \partial_2 = -\partial_t, \quad \partial_3 = \partial_t - \partial_y.$$

Let $\mathbf{r} = (x, y)$ stands for the space-like vector. The standard Lagrange transform relating Lagrangians and Hamiltonians gives

$$(15) \quad \mathcal{H} = \int d^2\mathbf{r} (A_a^* \partial_y A_a + A_c^* \partial_x A_c + V),$$

so that the equations of motion (7) and (9) are

$$(16) \quad \frac{d}{dt} A_a(\mathbf{r}, t) = \left\{ \mathcal{H}, A_a(\mathbf{r}, t) \right\}, \quad \text{etc.}$$

where the Poisson brackets are defined by

$$(17) \quad \left\{ A_v^*(\mathbf{r}', t), A_v(\mathbf{r}, t) \right\} = \delta(\mathbf{r}' - \mathbf{r}), \quad v = a, b, c.$$

Any other same-time bracket is zero.

Relations (10) and (14) are just one of many possible conventions for the field notations and space-time separation. All such conventions in the continuous case are equivalent. The choice (10) and the signs in (14) at this moment may be considered as an odd decision of the author.

2. THE DISCRETIZATION

Now we proceed to the discrete analogue of the standard and modified three-wave equations (7,9). The straightforward discretization of three-dimensional space gives the cubic lattice

$$(18) \quad \mathbf{x} = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + x_3 \mathbf{e}_3, \quad x_i \in \mathbb{R} \quad \mapsto \quad \mathbf{n} = n_1 \mathbf{e}_1 + n_2 \mathbf{e}_2 + n_3 \mathbf{e}_3, \quad n_i \in \mathbb{Z}$$

The discrete analogue of derivative is the difference derivative,

$$(19) \quad \partial_i \mapsto \Delta_i, \quad \Delta_i \phi(\mathbf{n}) \stackrel{\text{def}}{=} \phi(\mathbf{n} + \mathbf{e}_i) - \phi(\mathbf{n}).$$

The point is that we apply the straightforward discretization to the linear problems (6,8), discrete equations of motion must appear as the consistency conditions.

2.1. Standard Three-wave equations. The discrete linear problem corresponding to (6),

$$(20) \quad \Delta_i \psi_{j,\mathbf{n}} = A_{ij}(\mathbf{n}) \psi_{i,\mathbf{n}},$$

provides the discrete equations of motion

$$(21) \quad A_{jk}(\mathbf{n} + \mathbf{e}_i) = \frac{A_{jk}(\mathbf{n}) + A_{ji}(\mathbf{n})A_{ik}(\mathbf{n})}{1 - A_{ji}(\mathbf{n})A_{ij}(\mathbf{n})}.$$

2.2. Modified Three-wave equations. The discrete linear problem corresponding to (8),

$$(22) \quad \phi_{\mathbf{n}+\mathbf{e}_i+\mathbf{e}_j} - Q_{ji}(\mathbf{n})\phi_{\mathbf{n}+\mathbf{e}_i} - Q_{ij}(\mathbf{n})\phi_{\mathbf{n}+\mathbf{e}_j} + Q_{ij}(\mathbf{n})Q_{ji}(\mathbf{n})\phi_{\mathbf{n}} = 0,$$

where

$$(23) \quad Q_{ij}(\mathbf{n}) = 1 + A_{ij}(\mathbf{n}), \quad \text{etc.},$$

provides

$$(24) \quad Q_{jk}(\mathbf{n} + \mathbf{e}_i) = \frac{Q_{ji}(\mathbf{n})Q_{ik}(\mathbf{n}) + Q_{ij}(\mathbf{n})Q_{jk}(\mathbf{n}) - Q_{ij}(\mathbf{n})Q_{ji}(\mathbf{n})}{Q_{ik}(\mathbf{n})}.$$

3. CONSTANT TIME SURFACE AND EVOLUTION

Discrete equations of motion (21,24) evidently define a sort of one-step evolution: the fields at point $\mathbf{n} + \mathbf{e}_i$ are expressed in the terms of the fields at point $\mathbf{n}$. However, an explicit form of a space-like discrete surface and detailed definition of a discrete time corresponding to our choice (14) needs some discussion.

The key point is the geometrical structure of our discretization and the geometrical structure of the linear problems (20,22).

Let vector $\mathbf{n}$ in (18) stands for the vertex of the cubic lattice. The cubic lattice consists of *vertices* $\mathbf{n}$, *edges* $(\mathbf{n}, \mathbf{n} + \mathbf{e}_i)$, and *faces* $(\mathbf{n}, \mathbf{n} + \mathbf{e}_i, \mathbf{n} + \mathbf{e}_j, \mathbf{n} + \mathbf{e}_i + \mathbf{e}_j)$. The linear variable $\psi_{i,\mathbf{n}}$ of (20) should be associated with $(\mathbf{n}, \mathbf{n} + \mathbf{e}_i)$ -edge, the linear variable $\phi_{\mathbf{n}}$ should be associated with $\mathbf{n}$ -vertex, and the fields $A_{ij}(\mathbf{n})$, $A_{ji}(\mathbf{n})$ should be associated with $(\mathbf{n}, \mathbf{n} + \mathbf{e}_i, \mathbf{n} + \mathbf{e}_j, \mathbf{n} + \mathbf{e}_i + \mathbf{e}_j)$ -face. This is justified by the structure of the linear equations, see Fig. 1 and Fig. 2. In what follows, symbol $\mathcal{A}_{v,\mathbf{n}}$ stands for the collection of fields on $(v, \mathbf{n})$ -th face.




FIGURE 1. Graphical representation of the elements of the linear equations (20): linear variables $\psi_{i,\mathbf{n}}$ are associated with the edges of cubic lattice, the fields $\mathcal{A}_{a,\mathbf{n}} = (A_{12}(\mathbf{n}), A_{21}(\mathbf{n}))$ are associated with the face of cubic lattice.




FIGURE 2. Graphical representation of the elements of the linear equation (22): linear variables $\phi_{\mathbf{n}}$ are associated with the vertex of cubic lattice, the fields $\mathcal{A}_{a,\mathbf{n}} = (Q_{12}(\mathbf{n}), Q_{21}(\mathbf{n}))$ are associated with the face of cubic lattice.

Take now up the auxiliary linear problem (20) and more detailed derivation of (21). There are two ways to express $\psi_{1,\mathbf{n}+\mathbf{e}_2+\mathbf{e}_3}$, $\psi_{2,\mathbf{n}+\mathbf{e}_3}$, $\psi_{3,\mathbf{n}+\mathbf{e}_1+\mathbf{e}_2}$ in the terms of $\psi_{1,\mathbf{n}}$, $\psi_{2,\mathbf{n}+\mathbf{e}_1}$, $\psi_{3,\mathbf{n}}$. The one way is to use the six relations

$$(25) \quad \begin{aligned} \Delta_3 \psi_{1,\mathbf{n}+\mathbf{e}_2} &= A_{31}(\mathbf{n} + \mathbf{e}_2) \psi_{3,\mathbf{n}+\mathbf{e}_2}, & \Delta_3 \psi_{2,\mathbf{n}} &= A_{32}(\mathbf{n}) \psi_{3,\mathbf{n}}, & \Delta_2 \psi_{1,\mathbf{n}} &= A_{21}(\mathbf{n}) \psi_{2,\mathbf{n}}, \\ \Delta_1 \psi_{3,\mathbf{n}+\mathbf{e}_2} &= A_{13}(\mathbf{n} + \mathbf{e}_2) \psi_{1,\mathbf{n}+\mathbf{e}_2}, & \Delta_2 \psi_{3,\mathbf{n}} &= A_{23}(\mathbf{n}) \psi_{2,\mathbf{n}}, & \Delta_1 \psi_{2,\mathbf{n}} &= A_{12}(\mathbf{n}) \psi_{1,\mathbf{n}}. \end{aligned}$$

The second way is to use the six other relations

$$(26) \quad \begin{aligned} \Delta_3 \psi_{2,\mathbf{n}+\mathbf{e}_1} &= A_{32}(\mathbf{n} + \mathbf{e}_1) \psi_{3,\mathbf{n}+\mathbf{e}_1}, & \Delta_2 \psi_{1,\mathbf{n}+\mathbf{e}_3} &= A_{21}(\mathbf{n} + \mathbf{e}_3) \psi_{2,\mathbf{n}+\mathbf{e}_3}, & \Delta_3 \psi_{1,\mathbf{n}} &= A_{31}(\mathbf{n}) \psi_{3,\mathbf{n}}, \\ \Delta_2 \psi_{3,\mathbf{n}+\mathbf{e}_1} &= A_{23}(\mathbf{n} + \mathbf{e}_1) \psi_{2,\mathbf{n}+\mathbf{e}_1}, & \Delta_1 \psi_{2,\mathbf{n}+\mathbf{e}_3} &= A_{12}(\mathbf{n} + \mathbf{e}_3) \psi_{1,\mathbf{n}+\mathbf{e}_3}, & \Delta_1 \psi_{3,\mathbf{n}} &= A_{13}(\mathbf{n}) \psi_{1,\mathbf{n}}. \end{aligned}$$

Graphical representation for (25) and (26) is the left and right hand sides of Fig. 3. The collection of relations (25) and (26) correspond to the faces of the cube $(\mathbf{n}, \mathbf{n} + \mathbf{e}_i, \mathbf{n} + \mathbf{e}_i + \mathbf{e}_j, \mathbf{n} + \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3)$ – the discrete space-time consistency is the consistency around the cube.

The way to introduce the discrete time is to identify the discrete space-time fields in the left hand side of Fig. 3 with the discrete time t , and identify the discrete space-time fields in the right hand side of Fig. 3 with the discrete time $t + 1$. This exactly corresponds to (14):

$$(27) \quad \mathbf{e}_1 = \mathbf{e}_t - \mathbf{e}_x, \quad \mathbf{e}_2 = -\mathbf{e}_t, \quad \mathbf{e}_3 = \mathbf{e}_t - \mathbf{e}_y,$$




FIGURE 3. Consistency around the cube for the linear problem of Fig 1. Here $\psi_1 = \psi_{1,\mathbf{n}}$, $\psi_{1,2} = \psi_{1,\mathbf{n}+\mathbf{e}_2}$, $\psi_{1,23} = \psi_{1,\mathbf{n}+\mathbf{e}_2+\mathbf{e}_3}$, etc. The left hand side corresponds to Eqs. (25), the right hand side corresponds to Eqs. (26); $\mathcal{A}_{a,\mathbf{n}} = (A_{12}(\mathbf{n}), A_{21}(\mathbf{n}))$, etc., according to (10).

and therefore

$$(28) \quad \mathbf{n} = t\mathbf{e}_t + x\mathbf{e}_x + y\mathbf{e}_y : \quad x = -n_1, \quad y = -n_3, \quad t = n_1 - n_2 + n_3.$$

The following table gives the correspondence between the initial $A_{ij}(\mathbf{n})$ - notations and the space-time notations, convention (10) is taken into account:

$$(29) \quad \begin{aligned} A_{12}(\mathbf{n}) &= A_a(x, y, t) = A_a, & A_{21}(\mathbf{n}) &= A_a^*(x, y, t) = A_a^*, \\ A_{13}(\mathbf{n} + \mathbf{e}_2) &= A_b(x, y, t) = A_b, & A_{31}(\mathbf{n} + \mathbf{e}_2) &= A_b^*(x, y, t) = A_b^*, \\ A_{23}(\mathbf{n}) &= A_c(x, y, t) = A_c, & A_{32}(\mathbf{n}) &= A_c^*(x, y, t) = A_c^*. \end{aligned}$$

Here the third columns are the shortened notations. The right hand side of Fig. 3 implies thus

$$(30) \quad \begin{aligned} A_{12}(\mathbf{n} + \mathbf{e}_3) &= A_a(x, y - 1, t + 1) = \overline{A}_a, & A_{21}(\mathbf{n} + \mathbf{e}_3) &= A_a^*(x, y - 1, t + 1) = \overline{A}_a^*, \\ A_{13}(\mathbf{n}) &= A_b(x, y, t + 1) = \overline{A}_b, & A_{31}(\mathbf{n}) &= A_b^*(x, y, t) = \overline{A}_b^*, \\ A_{23}(\mathbf{n} + \mathbf{e}_1) &= A_c(x - 1, y, t + 1) = \overline{A}_c, & A_{32}(\mathbf{n} + \mathbf{e}_1) &= A_c^*(x - 1, y, t + 1) = \overline{A}_c^*, \end{aligned}$$

Here the third columns are the shortened notations as well. The consistency condition of (25) and (26), completely equivalent to Eq. (21), may be rewritten in shortened notations as

$$(31) \quad \left\{ \begin{aligned} \overline{A}_a^* &= A_a^* + A_b^* A_c, & \overline{A}_a &= \frac{A_a(1 - A_c A_c^*) + A_b A_c^*(1 - A_a A_a^*)}{1 - \overline{A}_b \overline{A}_b^*}, \\ \overline{A}_b^* &= A_b^*(1 - A_c A_c^*) - A_a^* A_c^*, & \overline{A}_b &= A_b(1 - A_a A_a^*) - A_a A_c, \\ \overline{A}_c^* &= \frac{A_c^*(1 - A_a A_a^*) + A_a A_b^*(1 - A_c A_c^*)}{1 - \overline{A}_b \overline{A}_c^*}, & \overline{A}_c &= A_c + A_a^* A_b. \end{aligned} \right.$$

In the absolutely similar way one may consider the discrete linear problem of Fig. 2. Corresponding picture is Fig. 4.




FIGURE 4. Consistency around the cube for the linear problem of Fig 2. Here $\phi = \phi_{\mathbf{n}}$, $\phi_1 = \psi_{\mathbf{n}+\mathbf{e}_1}$, $\phi_{12} = \phi_{\mathbf{n}+\mathbf{e}_1+\mathbf{e}_2}$, etc. Convention (10) is taken into account, $\mathcal{A}_{a,\mathbf{n}} = (Q_{12}(\mathbf{n}), Q_{21}(\mathbf{n}))$, etc.

Introducing the same-time variables for the left hand side of Fig. 4 analogously to (29),

$$(32) \quad \begin{aligned} Q_{12}(\mathbf{n}) &= u_a(x, y, t) = u_a, & Q_{21}(\mathbf{n}) &= w_a(x, y, t) = w_a, \\ Q_{13}(\mathbf{n} + \mathbf{e}_2) &= u_b(x, y, t) = u_b, & Q_{31}(\mathbf{n} + \mathbf{e}_2) &= w_b(x, y, t) = w_b, \\ Q_{23}(\mathbf{n}) &= u_c(x, y, t) = u_c, & Q_{32}(\mathbf{n}) &= w_c(x, y, t) = w_c, \end{aligned}$$

and for right hand side of Fig. 4 analogously to (30),

$$(33) \quad \begin{aligned} Q_{12}(\mathbf{n} + \mathbf{e}_3) &= u_a(x, y - 1, t + 1) = \bar{u}_a, & Q_{21}(\mathbf{n} + \mathbf{e}_3) &= w_a(x, y - 1, t + 1) = \bar{w}_a, \\ Q_{13}(\mathbf{n}) &= u_b(x, y, t + 1) = \bar{u}_b, & Q_{31}(\mathbf{n}) &= w_b(x, y, t + 1) = \bar{w}_b, \\ Q_{23}(\mathbf{n} + \mathbf{e}_1) &= u_c(x - 1, y, t + 1) = \bar{u}_c, & Q_{32}(\mathbf{n} + \mathbf{e}_1) &= w_c(x - 1, y, t + 1) = \bar{w}_c, \end{aligned}$$

we get the consistency condition in the shortened notations:

$$(34) \quad \left\{ \begin{aligned} \bar{u}_a &= \frac{u_a w_b + u_b w_c - u_b w_b}{w_c}, & \bar{w}_a &= \frac{w_a w_b u_c}{u_c w_c + w_a w_b - w_a w_c}, \\ \bar{u}_b &= \frac{u_a w_a + u_b u_c - u_a u_c}{w_a}, & \bar{w}_b &= \frac{u_c w_c + w_a w_b - w_a w_c}{u_c}, \\ \bar{u}_c &= \frac{w_a u_b u_c}{u_a w_a + u_b u_c - u_a u_c}, & \bar{w}_c &= \frac{u_a w_b + u_b w_c - u_b w_b}{u_a}. \end{aligned} \right.$$

These relations are completely equivalent to Eq. (24).

The derivation of the equations of motion as the consistency of linear problem around the cube is similar to the consistency approach to the integrable equations on quad-graphs [4]. The consistency around the cube for two-dimensional quad-graph equations is an analogue of the Yang-Baxter equation – variables in the left and right hand sides of two hexagons of Fig. 3 are the same. The consistency in this paper implies the different variables in the left and right hand sides, it defines a map. Therefore it is an analogue of the *local Yang-Baxter* equation or the *tetrahedral Zamolodchikov algebra* [5, 6, 7], see next section.

Equations (31) and (34) express the discrete space-time fields $\{\mathcal{A}_v(x, y, t+1) : v = a, b, c; x, y \in \mathbb{Z}\}$ in the terms of the fields $\{\mathcal{A}_v(x, y, t) : v = a, b, c; x, y \in \mathbb{Z}\}$. The space-like discrete surface is the collection of hexagons of Fig. 3 for all $x, y \in \mathbb{Z}$ and fixed t . This is the *honeycomb lattice*. Equations (31) and (34) therefore define the discrete time evolutions of the fields situated at the faces of the honeycomb lattices.

Often in the literature the dual lattices are used. The fields are associated to the edges of dual three-dimensional lattice and to the vertices of its section – dual two-dimensional lattice. The lattice dual to the honeycomb one is called the *kagome lattice*.

4. LINEAR PROBLEM AS THE ZERO CURVATURE REPRESENTATION

Equations (31) and (34) are identically equivalent to Eqs. (21) and (24) correspondingly, but look more complicated since we reverse the time direction of $\mathbf{e}_2$ in Eq. (27). Our way to introduce time and space-like coordinates (14,27) is not yet motivated.

Equations (31) define the map of variables $\mathcal{A}_v = (A_v, A_v^*)$ to $\overline{\mathcal{A}}_v = (\overline{A}_v, \overline{A}_v^*)$, $v = a, b, c$. Analogously, equation (34) define the map of variables $\mathcal{A}_v = (u_v, w_v)$ to $\overline{\mathcal{A}}_v = (\overline{u}_v, \overline{w}_v)$, $v = a, b, c$. Denote both these maps by the symbol $\mathcal{R}_{abc}$. Formally, $\mathcal{R}_{abc}$ is the operator acting in the space of functions of $\mathcal{A}_a, \mathcal{A}_b, \mathcal{A}_c$:

$$(35) \quad \forall \Phi = \Phi(\mathcal{A}_a, \mathcal{A}_b, \mathcal{A}_c) \quad : \quad \left(\mathcal{R}_{abc} \circ \Phi \right) (\mathcal{A}_a, \mathcal{A}_b, \mathcal{A}_c) \stackrel{\text{def}}{=} \Phi(\overline{\mathcal{A}}_a, \overline{\mathcal{A}}_b, \overline{\mathcal{A}}_c) .$$

The point is that the maps (31) and (34) are two basic set-theoretical solutions of the tetrahedron equation:

$$(36) \quad \mathcal{R}_{abc} \mathcal{R}_{ade} \mathcal{R}_{bdf} \mathcal{R}_{cef} = \mathcal{R}_{cef} \mathcal{R}_{bdf} \mathcal{R}_{ade} \mathcal{R}_{abc} .$$

It may be verified straightforwardly, see the appendix.

Take up now the question: why the maps (31) and (34) satisfy the functional tetrahedron equation. The reason is that the auxiliary linear problems defining the maps of dynamical variables from the left hand sides of Figs. 3 and 4 to the right hand sides are the *correctly oriented* zero curvature representations of three dimensional integrable model in discrete space-time.

Let me demonstrate this statement for the map (31). The linear equations for the single face, see Fig. 1,

$$(37) \quad \psi_{1, \mathbf{n} + \mathbf{e}_2} = \psi_{1, \mathbf{n}} + A_a^* \psi_{2, \mathbf{n}} , \quad \psi_{2, \mathbf{n} + \mathbf{e}_1} = \psi_{2, \mathbf{n}} + A_a \psi_{1, \mathbf{n}} .$$

may be rewritten in the matrix form as

$$(38) \quad \begin{pmatrix} \psi_{1, \mathbf{n}} \\ \psi_{2, \mathbf{n} + \mathbf{e}_1} \end{pmatrix} = X[\mathcal{A}_a] \cdot \begin{pmatrix} \psi_{1, \mathbf{n} + \mathbf{e}_2} \\ \psi_{2, \mathbf{n}} \end{pmatrix}, \quad X[\mathcal{A}_a] = \begin{pmatrix} 1 & -A_a^* \\ A_a & 1 - A_a A_a^* \end{pmatrix} .$$

The linear equations for all three faces of the left hand side of Fig. 3 may be written as

$$(39) \quad \begin{pmatrix} \psi_1 \\ \psi_{2,1} \end{pmatrix} = X[\mathcal{A}_a] \begin{pmatrix} \psi_{1,2} \\ \psi_1 \end{pmatrix}, \quad \begin{pmatrix} \psi_{1,2} \\ \psi_{3,12} \end{pmatrix} = X[\mathcal{A}_b] \begin{pmatrix} \psi_{1,23} \\ \psi_{3,2} \end{pmatrix}, \quad \begin{pmatrix} \psi_2 \\ \psi_{3,2} \end{pmatrix} = X[\mathcal{A}_c] \begin{pmatrix} \psi_{2,3} \\ \psi_3 \end{pmatrix}$$

with the same matrix function $X[\mathcal{A}]$ (38). Iterating these matrix equations, we come to

$$(40) \quad \begin{pmatrix} \psi_1 \\ \psi_{2,1} \\ \psi_{3,12} \end{pmatrix} = X_{12}[\mathcal{A}_a]X_{13}[\mathcal{A}_b]X_{23}[\mathcal{A}_c] \begin{pmatrix} \psi_{1,23} \\ \psi_{2,3} \\ \psi_3 \end{pmatrix},$$

where $X_{ij}[\mathcal{A}_v]$ are the three by three block-diagonal matrices, X_{ij} coincides with (38) in (ij) -block and has the unity in complimentary block. For instance,

$$(41) \quad X_{12}[\mathcal{A}_a] = \begin{pmatrix} 1 & -A_a^* & 0 \\ A_a & 1 - A_a A_a^* & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Analogously, the right hand side of Fig. 3 provides

$$(42) \quad \begin{pmatrix} \psi_1 \\ \psi_{2,1} \\ \psi_{3,12} \end{pmatrix} = X_{23}[\overline{\mathcal{A}}_a]X_{13}[\overline{\mathcal{A}}_b]X_{12}[\overline{\mathcal{A}}_c] \begin{pmatrix} \psi_{1,23} \\ \psi_{2,3} \\ \psi_3 \end{pmatrix}.$$

Thus, the consistency of the linear problem around the cube is the *Korepanov* equation [8, 6]

$$(43) \quad X_{12}[\mathcal{A}_a]X_{13}[\mathcal{A}_b]X_{23}[\mathcal{A}_c] = X_{23}[\overline{\mathcal{A}}_c]X_{13}[\overline{\mathcal{A}}_b]X_{12}[\overline{\mathcal{A}}_a].$$

Using definition (35), we may rewrite the Korepanov equation in the form similar to tetrahedral Zamolodchikov algebra [7],

$$(44) \quad X_{12}[\mathcal{A}_a]X_{13}[\mathcal{A}_b]X_{23}[\mathcal{A}_c] = \mathcal{R}_{abc} \circ X_{23}[\mathcal{A}_c]X_{13}[\mathcal{A}_b]X_{12}[\mathcal{A}_a].$$

Tetrahedron equation (36) is the equivalence of decompositions of the uniquely defined map $\mathcal{A}_v \rightarrow \overline{\overline{\mathcal{A}}}_v$

$$(45) \quad X_{12}[\mathcal{A}_a]X_{13}[\mathcal{A}_b]X_{23}[\mathcal{A}_c]X_{14}[\mathcal{A}_d]X_{24}[\mathcal{A}_e]X_{34}[\mathcal{A}_f] = X_{34}[\overline{\overline{\mathcal{A}}}_f]X_{24}[\overline{\overline{\mathcal{A}}}_e]X_{23}[\overline{\overline{\mathcal{A}}}_c]X_{14}[\overline{\overline{\mathcal{A}}}_d]X_{13}[\overline{\overline{\mathcal{A}}}_b]X_{12}[\overline{\overline{\mathcal{A}}}_a]$$

into two different sequences of elementary maps.

The main difference between (43) and the local Yang-Baxter equation [5, 6] is that the numerical indices in (43,44) correspond to the components of the *tensor sum* of one-dimensional vector spaces ψ_i . The Korepanov equation comes from the linear problem directly, therefore it is genuine multi-dimensional generalization of the Lax representation.

There is no analogous matrix form for the zero curvature representation of the modified three-wave equations. One has to work directly with the sets of linear relations (22). Nevertheless, the orientation of the faces in Fig. 4 is correct, and a set-of-linear-relations analysis similar to Eq. (45) provides the set-theoretical proof of the corresponding tetrahedron equation [9, 10].

5. POISSON BRACKETS FOR THE FUNDAMENTAL MAPS

The maps (31) and (34) define the discrete-time evolution on the honeycomb lattice. Evolution is the Hamiltonian one if it preserves Poisson brackets.

One may verify it for the map (34), if

$$(46) \quad \left\{ u_v, w_v \right\} = u_v w_v, \quad v = a, b, c$$

and any other type bracket for u_v, w_v is zero, then

$$(47) \quad \left\{ \overline{u}_v, \overline{w}_v \right\} = \overline{u}_v \overline{w}_v, \quad v = a, b, c$$

and any other type bracket for $\overline{u}_v, \overline{w}_v$ is zero. Therefore, (34) is the canonical map [9]. Restoring the space structure according to (32,33), we come to the whole system of same-time Poisson brackets conserved by the evolution:

$$(48) \quad \left\{ u_v(x, y, t), w_{v'}(x', y', t) \right\} = \delta_{v,v'} \delta_{x,x'} \delta_{y,y'} u_v(x, y, t) w_{v'}(x', y', t),$$

any other type same-time bracket is zero. Poisson algebra with such structure of delta-symbols is called the *ultra-local* one.

The Poisson brackets for the evolution (31) are not ultra-local. To get the ultra-locality, we need to modify the discrete linear problem (38). Let there

$$(49) \quad X[\mathcal{A}] = \begin{pmatrix} K & -A^* \\ A & K \end{pmatrix}, \quad \mathcal{A} = (A, A^*, K), \quad K^2 = 1 - AA^*.$$

The modified map comes from the Korepanov equation (43):

$$(50) \quad \begin{cases} \overline{A}_a^* = \overline{K}_b^{-1} (K_c A_a^* + K_a A_b^* A_c), & \overline{A}_a = \overline{K}_b^{-1} (K_c A_a + K_a A_b A_c^*), \\ \overline{A}_b^* = K_a K_c A_b^* - A_a^* A_c^*, & \overline{A}_b = K_a K_c A_b - A_a A_c, \\ \overline{A}_c^* = \overline{K}_b^{-1} (K_a A_c^* + K_c A_a A_b^*), & \overline{A}_c = \overline{K}_b^{-1} (K_a A_c + K_c A_a^* A_b). \end{cases}$$

By definition (49), $\overline{K}_v^2 = 1 - \overline{A}_v \overline{A}_v^*$. Additional property of this map is

$$(51) \quad \overline{K}_a \overline{K}_b = K_a K_b, \quad \overline{K}_b \overline{K}_c = K_b K_c.$$

The map (50) satisfies the functional tetrahedron equation and preserves the ultra-local Poisson brackets [11]

$$(52) \quad \left\{ A_v^*, A_v \right\} = K_a^2, \quad \left\{ K_v, A_v \right\} = -\frac{1}{2} K_v A_v, \quad \left\{ A_v^*, K_v \right\} = -\frac{1}{2} K_v A_v^*,$$

any other type bracket is zero.

The principal advantage of the Poisson structure is that it allows one to define the lattice actions with the help of functions generating the canonical transformations. This subject is technically complicated, we postpone it for future publications.

6. QUANTIZATION

The Poisson algebra (52) is the quasi-classical limit of q -oscillator algebra [12]

$$(53) \quad \mathbf{A}\mathbf{A}^\dagger = 1 - q^{2\mathbf{N}+2}, \quad \mathbf{A}^\dagger\mathbf{A} = 1 - q^{2\mathbf{N}}, \quad \mathbf{A}q^{\mathbf{N}} = q^{\mathbf{N}+1}\mathbf{A}, \quad \mathbf{A}^\dagger q^{\mathbf{N}} = q^{\mathbf{N}-1}\mathbf{A}^\dagger.$$

The Poisson bracket is the limit of commutator when $q^2 = e^{-\hbar} \rightarrow 1$, $\mathbf{A} \rightarrow A$, $\mathbf{A}^\dagger \rightarrow A^*$ and $q^{\mathbf{N}} \rightarrow K$.

The Poisson algebra $\{u, w\} = uw$ (46) is the quasi-classical limit of the Weyl algebra [13]

$$(54) \quad \mathbf{u}\mathbf{w} = q^2\mathbf{w}\mathbf{u}$$

with $\mathbf{u} \rightarrow u$, $\mathbf{w} \rightarrow w$ when $q \rightarrow 1$.

The whole algebra of observables, corresponding to the set of classical fields $\mathcal{A}_v(x, y)$ is thus the tensor power of the local q -oscillator algebra $\mathcal{A} = (\mathbf{A}, \mathbf{A}^\dagger, q^{\mathbf{N}})$ for the quantized three-wave system; and the tensor power of the local Weyl algebra $\mathcal{A} = (\mathbf{u}, \mathbf{w})$ for the modified three-wave system. After the quantization, the indices a, b, c of the operators (or, more generally, the indices are (v, x, y) , $v = a, b, c$, $x, y \in \mathbb{Z}$) stand for components of the tensor power.

Quantum maps follow from the quantum linear problems. Quantized version of (38) is [14]

$$(55) \quad \begin{aligned} |\psi_{1,\mathbf{n}}\rangle &= \lambda_a q^{\mathbf{N}_a} |\psi_{1,\mathbf{n}+\mathbf{e}_2}\rangle - \mathbf{A}^\dagger_a |\psi_{2,\mathbf{n}}\rangle, \\ |\psi_{2,\mathbf{n}+\mathbf{e}_1}\rangle &= q^{-1} \lambda_a \mu_a \mathbf{A}_a |\psi_{1,\mathbf{n}+\mathbf{e}_2}\rangle + \mu_a q^{\mathbf{N}_a} |\psi_{2,\mathbf{n}}\rangle \end{aligned} \Leftrightarrow X[\mathcal{A}_a] = \begin{pmatrix} \lambda_a q^{\mathbf{N}_a} & -\mathbf{A}^\dagger_a \\ q^{-1} \lambda_a \mu_a \mathbf{A}_a & \mu_a q^{\mathbf{N}_a} \end{pmatrix}$$

Here we replace the linear variables ψ_i by vectors $|\psi_i\rangle$ from a formal right module of the whole algebra of observables. Parameters λ_a, μ_a are $\mathbb{C}$ -valued spectral parameters, we introduce them for the sake of completeness. The consistency of linear problem of Fig. 3 (equivalent to the quantum Korepanov equation (43)) may be solved with non-commutative coefficients, the answer is [11, 14]

$$(56) \quad \left\{ \begin{aligned} \overline{\mathbf{A}^\dagger}_a &= \frac{\lambda_c}{\lambda_b} q^{-\overline{\mathbf{N}}_b} (q^{\mathbf{N}_c} \mathbf{A}^\dagger_a + \frac{\lambda_a \mu_c}{q} q^{\mathbf{N}_a} \mathbf{A}^\dagger_b \mathbf{A}_c), & \overline{\mathbf{A}}_a &= \frac{\lambda_b}{\lambda_c} q^{-\overline{\mathbf{N}}_b} (q^{\mathbf{N}_c} \mathbf{A}_a + \frac{q}{\lambda_a \mu_c} q^{\mathbf{N}_a} \mathbf{A}_b \mathbf{A}^\dagger_c), \\ \overline{\mathbf{A}^\dagger}_b &= \lambda_a \mu_c q^{\mathbf{N}_a + \mathbf{N}_c} \mathbf{A}^\dagger_b - \mathbf{A}^\dagger_a \mathbf{A}^\dagger_c, & \overline{\mathbf{A}}_b &= \frac{q^2}{\lambda_a \mu_c} q^{\mathbf{N}_a + \mathbf{N}_c} \mathbf{A}_b - \mathbf{A}_a \mathbf{A}_c, \\ \overline{\mathbf{A}^\dagger}_c &= \frac{\mu_a}{\mu_b} q^{-\overline{\mathbf{N}}_b} (q^{\mathbf{N}_a} \mathbf{A}^\dagger_c + \frac{\lambda_a \mu_c}{q} q^{\mathbf{N}_c} \mathbf{A}_a \mathbf{A}^\dagger_b), & \overline{\mathbf{A}}_c &= \frac{\mu_b}{\mu_a} q^{-\overline{\mathbf{N}}_b} (q^{\mathbf{N}_a} \mathbf{A}_c + \frac{q}{\lambda_a \mu_c} q^{\mathbf{N}_c} \mathbf{A}^\dagger_a \mathbf{A}_b). \end{aligned} \right.$$

Here $q^{2\overline{\mathbf{N}}_b} = 1 - \overline{\mathbf{A}^\dagger}_b \overline{\mathbf{A}}_b$, in addition

$$(57) \quad q^{\overline{\mathbf{N}}_a + \overline{\mathbf{N}}_b} = q^{\mathbf{N}_a + \mathbf{N}_b}, \quad q^{\overline{\mathbf{N}}_b + \overline{\mathbf{N}}_c} = q^{\mathbf{N}_b + \mathbf{N}_c},$$

One may verify, the map (56) is the *automorphism* of the tensor cube of q -oscillator algebra (53).

Quantized version of the linear equation (22) for the modified three-wave system is [10]

$$(58) \quad \varkappa_a |\phi_{\mathbf{n}+\mathbf{e}_1+\mathbf{e}_2}\rangle - q \mathbf{w}_a |\phi_{\mathbf{n}+\mathbf{e}_1}\rangle - \mathbf{u}_a |\phi_{\mathbf{n}+\mathbf{e}_2}\rangle + \mathbf{u}_a \mathbf{w}_a |\phi_{\mathbf{n}}\rangle = 0.$$

Here $\varkappa_a$ is a $\mathbb{C}$ -valued spectral parameter. Solution of the consistency condition, see Fig. 4, gives

$$(59) \quad \begin{cases} \overline{\mathbf{u}}_a = \Lambda_2 \mathbf{w}_c^{-1}, & \overline{\mathbf{u}}_b = \Lambda_1 \mathbf{u}_c, & \overline{\mathbf{u}}_c = \mathbf{u}_b \Lambda_1^{-1}, \\ \overline{\mathbf{w}}_a = \mathbf{w}_b \Lambda_3^{-1}, & \overline{\mathbf{w}}_b = \Lambda_3 \mathbf{w}_a, & \overline{\mathbf{w}}_c = \Lambda_2 \mathbf{u}_a^{-1}, \end{cases}$$

where

$$\begin{aligned}
\Lambda_1 &= \mathbf{u}_a \mathbf{u}_c^{-1} - q \mathbf{u}_a \mathbf{w}_a^{-1} + \varkappa_a \mathbf{u}_b \mathbf{w}_a^{-1} , \\
\Lambda_2 &= \frac{\varkappa_a}{\varkappa_b} \mathbf{u}_b \mathbf{w}_c + \frac{\varkappa_c}{\varkappa_b} \mathbf{u}_a \mathbf{w}_b - q^{-1} \frac{\varkappa_a \varkappa_c}{\varkappa_b} \mathbf{u}_b \mathbf{w}_b , \\
\Lambda_3 &= \mathbf{w}_a^{-1} \mathbf{w}_c - q \mathbf{u}_c^{-1} \mathbf{w}_c + \varkappa_c \mathbf{u}_c^{-1} \mathbf{w}_b .
\end{aligned}
\tag{60}$$

One may verify, the map (59) is the automorphism of the tensor cube of the Weyl algebra (54).

Both automorphisms (56) and (59) satisfy the “functional” tetrahedron equation. In irreducible representations they are the internal automorphisms,

$$\mathcal{R}_{abc} \circ \Phi \equiv R_{abc} \Phi R_{abc}^{-1} .
\tag{61}$$

Corresponding operators R_{abc} satisfy the quantum (operator-valued) tetrahedron equations. Matrix elements of R_{abc} are functions of the spectral parameters λ_v, μ_v for (56) and $\varkappa_v$ for (59).

The local maps (56) and (59) define the evolution map on the honeycomb lattice via the identification

$$\begin{aligned}
\mathcal{A}_a &= \mathcal{A}_{a,x,y}(t) & \overline{\mathcal{A}}_a &= \mathcal{A}_{a,x,y-1}(t+1) \\
\mathcal{A}_b &= \mathcal{A}_{b,x,y}(t) & \rightarrow & \overline{\mathcal{A}}_b = \mathcal{A}_{b,x,y}(t+1) \\
\mathcal{A}_c &= \mathcal{A}_{c,x,y}(t) & \overline{\mathcal{A}}_c &= \mathcal{A}_{c,x-1,y}(t+1)
\end{aligned}
\tag{62}$$

in accordance with (29,30) and (32,33). The evolution map is the automorphism of the whole algebra of observables. Parameters λ_v, μ_v for the q -oscillator model and κ_v for the Weyl algebra model should be (x, y) -independent. Then, in proper representations, the evolution is the internal automorphism given by an evolution operator,

$$\Phi(t+1) = U \Phi(t) U^{-1} .
\tag{63}$$

Matrix elements of U may be constructed with the help of matrix elements of local R_{abc} .

Examples of irreducible representations providing a “good” quantum mechanics are the following. For the q -oscillator algebra it is the case of real $q^2 = e^{-\hbar}$, $0 < q < 1$, and the Fock space: the Fock vacuum is defined by $\mathbf{A}|0\rangle = \mathbf{N}|0\rangle = 0$. The dagger of $\mathbf{A}^\dagger$ stands for the Hermitian conjugation, $\mathbf{N}^\dagger = \mathbf{N}$. If in addition $\frac{\lambda_c}{\lambda_b}, \frac{\mu_a}{\mu_b}$ and $\frac{\lambda_a \mu_c}{q}$ in (56) are unitary parameters, then R_{abc} and U are the well defined unitary operators. Matrix elements of R_{abc} are given in [11]. Evaluation (4) of U^T in the framework of normal symbols gives the Feynman-type integral in the discrete space-time. In the quasi-classical limit $\hbar \rightarrow 0$ ($q^2 = e^{-\hbar}$), it may be shown

$$\langle A_{out}^* | U^T | A_{in} \rangle = \int \mathcal{D}A \mathcal{D}A^* e^{\frac{i}{\hbar} S[A, A^*]} ,
\tag{64}$$

where $S[A, A^*]$ is the lattice action mentioned in the previous section, and the measure of integration is as well the lattice one.

The proper quantum mechanical representation of the Weyl algebra is given by the modular dualization [15]. In addition to the given local Weyl pairs

$$\mathbf{u} = e^P , \quad \mathbf{w} = e^Q , \quad [Q, P] = i\hbar \Rightarrow q^2 = e^{-i\hbar} ,
\tag{65}$$

it is necessary to consider the dual local pairs

$$(66) \quad \mathbf{u}' = e^{\frac{2\pi}{\hbar}P}, \quad \mathbf{w}' = e^{\frac{2\pi}{\hbar}Q}, \quad q'^2 = e^{-i\frac{(2\pi)^2}{\hbar}}.$$

The sets of equations (59) and similar equations for dual pairs define the kernel of R_{abc} unambiguously. It is known, in this case R_{abc} and U are well defined unitary operators. The PQ -symbol of U^T in the quasi-classical limit $\hbar \rightarrow 0$ is

$$(67) \quad \langle P_{out}|U^T|Q_{in}\rangle = \int \mathcal{D}P\mathcal{D}Q e^{\frac{i}{\hbar}S[P,Q]}$$

where the measure and the action are the lattice ones.

7. CONCLUSION

The honeycomb lattice evolution map (62) defined with the help of local automorphisms (56) and (59), and the proper choice of the Hilbert space define unambiguously the evolution operator for the lattice approximation of corresponding quantum field theory. These quantum field theories are integrable since the underlying quantum auxiliary linear problems (55) and (58) provide the existence of the complete set of the integrals of motion [8, 16, 17, 18, 14].

The one-step evolution (62) is the discrete form of the Hamiltonian flow (15,16). The quantum lattice Hamiltonian $\mathbf{H}$ is defined by

$$(68) \quad U = e^{-\frac{i}{\hbar}\mathbf{H}\Delta t},$$

where $\Delta t = \Delta x = \Delta y$ is the lattice spacing parameter. In the continuous $\Delta t \rightarrow 0$ classical $q \rightarrow 1$ limit the quantum lattice Hamiltonian becomes exactly (15). However, on the lattice Δt is finite (in this paper we used the scale $\Delta t = 1$), and therefore $\mathbf{H}$ is not a polynomial in the algebra of observables. The evolution operator is the principal object of the field theory rather than a Hamiltonian.

The fundamental problem of the field theory is the calculation of the spectrum of evolution operator. This is the open question in three-dimensional models. Spectral equations for the evolution operators are not known yet (except for the two-dimensional limit of the three-wave system, [19]).

Let me conclude the paper with a discussion of the role of simplex equations in quantum field theory. The linear problem is the starting point of the integrability. The tetrahedron equation (36) and the Yang-Baxter equation for the two-dimensional models are the elementary consequences of linear problem as the zero curvature representation. Nevertheless, the simplex configurations may be considered as fragments of the space-time lattice, for instance

$$(69) \quad Z = \int \underbrace{d\phi'_1 d\phi'_2 d\phi'_3}_{\mathcal{D}\phi} \dots \underbrace{\langle \phi_1 \phi_2 | R_{12} | \phi'_1, \phi'_2 \rangle \langle \phi'_1, \phi_3 | R_{13} | \phi''_1, \phi'_3 \rangle \langle \phi'_2, \phi'_3 | R_{23} | \phi''_2, \phi''_3 \rangle}_{e^{\frac{i}{\hbar}S[\phi]}} \dots$$

Here for brevity we consider a two-dimensional theory and the triangle configuration. Such partition function corresponds to the Feynman integral (1) with the discrete measure definition (2) and some particular discretization. The Yang-Baxter equation (and the D -simplex equations in general) is the condition of Z -invariance: the partition function (69) does not depend on particular details of the

discretization, it is an invariant function of the boundary fields only. Thus the natural role of simplex equations in the lattice field theory is the conditions of self-consistent definition of the measure (2).

Acknowledgements I would like to thank all the participants of SIDE VII conference for fruitful discussions.

REFERENCES

- [1] V. E. Zakharov and S. V. Manakov, “Resonant interaction of wave packets in nonlinear media”, *JETP Lett.* **18** (1973) 243-245.
V. E. Zakharov and S. V. Manakov, “The theory of resonance interaction of wave packets in nonlinear media”, *Sov. Phys. JETP* **42** (1975) 842–850
- [2] D. J. Kaup, “The inverse scattering solution for the full three dimensional three-wave resonant interaction”, *Physica* **1D** (1980) 45-67
- [3] V. V. Mangazeev and S. M. Sergeev, “The continuous limit of the triple τ -function model”, *Theoretical and Mathematical Physics* **129** (2001) 317–326
- [4] A. I. Bobenko and Yu. B. Suris, “Integrable noncommutative equations on quad-graphs. The consistency approach”, *Lett. Math. Phys.* **61** (2002) 241–254
V. E. Adler, A. I. Bobenko and Yu. B. Suris, “Classification of integrable equations on quad-graphs. The consistency approach.” *Comm. Math. Phys.* **233** (2003) 513–543
- [5] J.-M. Maillet and F. Nijhoff, “Integrability for three dimensional lattice models”, *Phys. Lett.* **B224** 389-396
- [6] R. M. Kashaev, I. G. Korepanov and S. M. Sergeev, “Functional Tetrahedron Equation”, *Theoretical and Mathematical Physics* **117** (1998) 370-384
- [7] I. G. Korepanov, “Tetrahedral Zamolodchikov algebras corresponding to Baxter’s L -operators”, *J. Stat. Phys.* **71** (1993) 85-97
- [8] I. G. Korepanov, “Algebraic integrable dynamical systems, $2+1$ dimensional models on wholly discrete space-time, and inhomogeneous models on 2-dimensional statistical physics”, *Preprint* arXiv:solv-int/9506003
- [9] S. M. Sergeev, “3D symplectic map”, *Phys. Lett. A* **253** (1999) 145–150
- [10] S. M. Sergeev, “Quantum $2+1$ evolution model”, *J. Phys. A: Math. Gen.* **32** (1999) 5693–5714
- [11] V. V. Bazhanov and S. M. Sergeev, “Zamolodchikov’s tetrahedron Equation and Hidden Structure of Quantum Groups”, *J. Phys. A: Math. Gen.* **39** (2006) 3295-3310
- [12] E. V. Damaskinsky and P. P. Kulish, “Deformed oscillators and their applications”, *J. Soviet Math.* **62** (1992) 2963–2986
- [13] V. V. Bazhanov and N. Yu. Reshetikhin, “Remarks on the quantum dilogarithm”, *J. Phys. A* **28** (1995) 2217–2226
- [14] S. Sergeev, “Quantum curve in q -oscillator model”, *preprint* nlin.SI/0510048, accepted in *International Journal of Mathematics and Mathematical Sciences* (2006)
- [15] L. D. Faddeev, “Discrete Heisenberg-Weyl group and modular group”, *Lett. Math. Phys.* **34** (1995) 249–254
- [16] S. M. Sergeev, “Auxiliary transfer matrices for three-dimensional integrable models”, *Theoretical and Mathematical Physics* **124** (2000) 391–409
- [17] S. M. Sergeev, “Quantum matrices of the coefficients of a discrete linear problem”, (*Russian*) *Zap. Nauchn. Sem. S.-Petersburg. Otdel. Mat. Inst. Steklov. (POMI)* **269** No. 16 (2000) “Vopr. Kvant. Teor. Polya i Stat. Fiz.” 292–307, 370–371 Translation: “Coefficient Matrices of a Quantum Discrete Auxiliary Linear Problem”, *Journal of Mathematical Sciences* **115**(1) (2003) 2049-2057
- [18] S. Sergeev, “Integrability of q -oscillator lattice model”, *preprint Physics Letters A* **357** (2006) 417-419
- [19] S. Sergeev, “Evolution operators for quantum chains”, *Preprint* arXiv:nlin.SI/0607055

APPENDIX A. VERIFICATION OF THE FUNCTIONAL TETRAHEDRON EQUATIONS

The functional tetrahedron equation for the map (31) may be verified with the help of Maple 9.5 routine:

```
restart;

R:=proc(a,b,c,var) # x stands for A, y stands for A^*
  local X,Y;
  X[2]:= x[b]*(1-x[a]*y[a])-x[a]*x[c];
  Y[2]:= y[b]*(1-x[c]*y[c])-y[a]*y[c];
  X[1]:=(x[a]*(1-x[c]*y[c])+x[b]*y[c]*(1-x[a]*y[a]))/(1-X[2]*Y[2]);
  Y[1]:=y[a]+y[b]*x[c]; X[3]:=x[c]+y[a]*x[b];
  Y[3]:=(y[c]*(1-x[a]*y[a])+x[a]*y[b]*(1-x[c]*y[c]))/(1-X[2]*Y[2]);
  simplify(subs([x[a]=X[1],y[a]=Y[1],x[b]=X[2],y[b]=Y[2],x[c]=X[3],y[c]=Y[3]],var))
end;
```

```
TE:=var->R(1,2,3,R(1,4,5,R(2,4,6,R(3,5,6,var))))
  -R(3,5,6,R(2,4,6,R(1,4,5,R(1,2,3,var))));
```

```
for k from 1 to 6 do TE(x[k]); TE(y[k]); od;
```

Verification of the tetrahedron equation for the map (34) is

```
restart;

R:=proc(a,b,c,var)
  local U,W;
  U[1]:=(u[a]*w[b]+u[b]*w[c]-u[b]*w[b])/w[c];
  W[1]:=w[a]*w[b]*u[c]/(u[c]*w[c]+w[a]*w[b]-w[a]*w[c]);
  U[2]:=(u[a]*w[a]+u[b]*u[c]-u[a]*u[c])/w[a];
  W[2]:=(u[c]*w[c]+w[a]*w[b]-w[a]*w[c])/u[c];
  U[3]:=w[a]*u[b]*u[c]/(u[a]*w[a]+u[b]*u[c]-u[a]*u[c]);
  W[3]:=(u[a]*w[b]+u[b]*w[c]-u[b]*w[b])/u[a];
  simplify(subs([u[a]=U[1],w[a]=W[1],u[b]=U[2],w[b]=W[2],u[c]=U[3],w[c]=W[3]],var))
end;
```

```
TE:=var->R(1,2,3,R(1,4,5,R(2,4,6,R(3,5,6,var))))
  -R(3,5,6,R(2,4,6,R(1,4,5,R(1,2,3,var))));
```

```
for k from 1 to 6 do TE(u[k]); TE(w[k]); od;
```

DEPARTMENT OF THEORETICAL PHYSICS (RESEARCH SCHOOL OF PHYSICAL SCIENCES AND ENGINEERING) & MATHEMATICAL SCIENCE INSTITUTE, CANBERRA ACT 0200, AUSTRALIA

E-mail address: `sergey.sergeev@anu.edu.au`